

Supplementary Table 6. Comparative Analysis of Endoscopic Phenotype between Patients with and without Pouch Failure (n = 50)

Variable	Category	DLI (-) (n = 45)	DLI (+) (n = 5)	P-value ^a
Normal	Yes	0	0	-
	No	45 (100)	5 (100)	
	NA	0	0	
Afferent limb involvement	Yes	10 (22.2)	1 (20)	0.333
	No	22 (48.9)	0	
	NA	13 (28.9)	4 (80)	
Inlet limb involvement	Yes	15 (33.3)	2 (40)	0.565
	No	22 (48.9)	1 (20)	
	NA	8 (17.8)	2 (40)	
Diffuse inflammation	Yes	7 (15.6)	4 (80)	0.006
	No	38 (84.4)	1 (20)	
	NA	0	0	
Focal inflammation	Yes	38 (84.4)	1 (20)	0.006
	No	7 (15.6)	4 (80)	
	NA	0	0	
Cuffitis	Yes	32 (71.1)	4 (80)	1.000
	No	13 (28.9)	1 (20)	
	NA	0	0	
Pouch fistula	Yes	2 (4.4)	5 (100)	<0.001
	No	43 (95.6)	0	
	NA	0	0	

Values are presented as number (%).

^aFisher exact test.

DLI, diverting loop ileostomy.