

Supplementary Table 5. WPAI and Annual Indirect Cost among Treated UC Patients Who Were Employed and Aged 18–64 Years

Continuous variable	No.	Mean	Median	SD	IQR	Range
WPAI ^a						
Absenteeism %	146	5.6	0.0	16.7	0–0	0.0–100.0
Presenteeism %	162	28.5	20.0	28.2	0–50	0.0–100.0
Overall work impairment %	145	30.3	20.0	30.0	0–50	0.0–100.0
Activity impairment %	164	27.6	20.0	28.1	0–50	0.0–100.0
Annual indirect cost from loss in work productivity (in million, JPY) ^b						
Absenteeism cost	146	0.2	0.0	0.6	0.0–0.0	0.0–3.9
Presenteeism cost	162	1.1	0.7	1.1	0–2.0	0.0–5.0
Indirect cost	145	1.1	0.8	1.2	0–1.9	0.0–5.0

^aThe sample sizes pertaining to absenteeism, presenteeism, and overall work impairment rely on respondents who provided responses to relevant questionnaires within the employed patient population.

^bJapanese Yen (¥1,000 = \$6.44).

WPAI, work productivity and activity impairment; UC, ulcerative colitis; SD, standard deviation; IQR, interquartile range.